

view. Navigation software is also progressing; luminal "fly through" navigation can be automated and collision avoidance techniques used, with simultaneous antegrade and retrograde viewing.

At present, full colonic cleansing is mandatory for an acceptable examination, but workers are already investigating the possibility of using oral contrast medium with a view to tagging faecal residue so that it can be differentiated from pathological features and possibly digitally excluded from the image. Furthermore, because virtual colonoscopy generates images of the colonic wall, it is possible to apply algorithms that automatically detect and label regions where it is thickened, alerting the radiologist to the possible presence of polyps. Perhaps flat adenomas can be detected this way.

The development and refinement of all of these techniques are likely to have tremendous impact on the speed and accuracy of virtual colonoscopy and its interpretation in the near future. The performance characteristics of virtual colonoscopy are certain to improve with improvements in hardware and software innovation and radiologists' experience. With regard to its potential as a screening tool, its performance in real populations and its cost, patient acceptability, and availability will need to be determined.

Competing interests: None declared.

- 1 Hounsfield GN. Computerized transverse axial scanning (tomography): Part 1. Description of the system. *Br J Radiol* 1973;46:1016-22.

- 2 Kalender WA, Seissler W, Klotz E, Vock P. Spiral volumetric CT with single breath-hold technique, continuous transport and continuous scanner rotation. *Radiology* 1990;176:181-3.
- 3 Fink M, Freeman AH, Dixon AK, Coni NK. Computed tomography of the colon in elderly people. *BMJ* 1994;308:101.
- 4 Amin Z, Poulos PB, Lees WR. Spiral CT pneumocolon for suspected colonic neoplasms. *Clin Radiol* 1996;51:56-61.
- 5 Vining DJ, Gelfand DW, Bechtold RE, Scharding ES, Grishaw EK, Shifrin RY, et al. Technical feasibility of colon imaging with helical CT and virtual reality [abstract]. *AJR Am J Roentgenol* 1994;162(S):104.
- 6 Hara AK, Johnson CD, Reed JE, Ahlquist DA, Nelson H, MacCarty RL, et al. Detection of colorectal polyps with CT colography: initial assessment of sensitivity and specificity. *Radiology* 1997;205:59-65.
- 7 Hara AK, Johnson CD, Reed JE, Ahlquist DA, Nelson H, Harmsen WS, et al. Computed tomographic colography for polyp detection: early comparison against barium enema. *Gastroenterology* 1997;112:A575.
- 8 Rex DK, Vining D, Kopecky K. Screening for polyps using spiral CT with and without virtual colonoscopy [abstract]. *Gastrointest Endosc* 1997;45:AB116.
- 9 Kay CL, Kulling D, Evandelou H, Hawes RH, Young JW, Cotton PB. CT-based virtual colonoscopy—correlation with colonoscopy in the detection of space-occupying lesions of the colon [abstract]. *Radiology* 1997;205:195.
- 10 Dachman AH, Kuniyoshi JK, Boyle CM, Samara Y, Hoffmann KR, Rubin DT, et al. CT colonography with three-dimensional problem solving for detection of colonic polyps. *AJR Am J Roentgenol* 1998;171:989-95.
- 11 Royster AP, Fenlon HM, Clarke PD, Nunes DP, Ferrucci JT. CT colonoscopy of colorectal neoplasms: two-dimensional and three-dimensional virtual reality techniques with colonoscopic correlation. *AJR Am J Roentgenol* 1997;169:1237-42.
- 12 Fenlon HM, McAneeny DB, Nunes DP, Clarke PD, Ferrucci JT. Occlusive colon carcinoma: virtual colonoscopy in the preoperative evaluation of the proximal colon. *Radiology* 1999;210:423-8.
- 13 Fenlon HM, Nunes DP, Clarke PD, Ferrucci JT. Colorectal neoplasm detection using virtual colonoscopy: a feasibility study. *Gut* 1998;43:806-11.
- 14 Lubolt W, Steiner P, Bauerfeind P, Pelkonen P, Debatin JF. Detection of mass lesions with MR colonography: preliminary report. *Radiology* 1998;207:59-65.
- 15 Fenlon HM, Ferrucci JT. Virtual colonoscopy. What will the issues be? *AJR Am J Roentgenol* 1997;169:453-8.

## Lesson of the week

### Oestrogen and calcium homeostasis in women with hypoparathyroidism

Janet McIlroy, Frances Dryburgh, John Hinnie, Rosemary Dargie, Alaa Al-Rawi

Oestrogen status can affect the requirement for vitamin D in women with hypoparathyroidism. Permanent hypoparathyroidism occurs in 0.2%–0.3% of patients who undergo thyroid surgery<sup>1</sup>; it arises less frequently as an inherited or as an autoimmune disease. The condition is treated with vitamin D analogue drugs, doses of which have to be titrated against the serum calcium concentration while avoiding hypercalciuria. The vitamin D requirements in women with hypoparathyroidism can change if their oestrogen status alters. An awareness of this can avoid hypercalcaemia.

#### Case reports

##### Case 1

A 54 year old woman had undergone thyroidectomy for Graves' disease 34 years previously and had been treated with vitamin D since that time. She had remained euthyroid, and her serum calcium concentration had been satisfactory and stable for several years on 1 $\alpha$ -hydroxycholecalciferol treatment (1 g/day). The patient's parathyroid hormone concentration was below the level of detection, although her calcitonin concentration was measurable (32 ng/l, reference range < 45 ng/l). Two months after stopping hormone replacement therapy she developed symptoms of hypercalcaemia—anorexia,

nausea, abdominal pain, constipation, and weight loss of 9 kg. Hypercalcaemia was confirmed biochemically; her calcium concentration, adjusted for albumin, was 3.5 mmol/l. Her dose of 1 $\alpha$ -hydroxycholecalciferol, calcium intake, and compliance with treatment were unchanged. She had been on hormone replacement therapy for 8 years—a cyclical regimen had been prescribed initially, and thereafter a continuous combined preparation. After 1 $\alpha$ -hydroxycholecalciferol was stopped, her calcium concentration returned to normal. Her serum calcium concentration subsequently remained within the reference range on a reduced dose of 1 $\alpha$ -hydroxycholecalciferol (0.25 g/day).

##### Case 2

A 52 year old woman with idiopathic hypoparathyroidism had been treated for 30 years with vitamin D analogues. For many years she had been taking 1 $\alpha$ -hydroxycholecalciferol (3 g/day). At annual review she was hypercalcaemic (calcium concentration adjusted for albumin, 3.1 mmol/l) and had anorexia, nausea, and weight loss of 12 kg. Before this her serum calcium concentration had consistently been within the reference range. There had been no change in her

Department of Clinical Biochemistry, Macewen Building, Glasgow Royal Infirmary, Glasgow. G4 0SF  
Janet McIlroy specialist registrar  
Frances Dryburgh consultant

continued over

BMJ 1999;319:1252–3

calcium intake or in compliance with treatment. Her calcium concentration returned to normal values after the dosage of 1 $\alpha$ -hydroxycholecalciferol was reduced to 1 g on alternate days. Further inquiry showed that her last menstrual period had been 3 months before the discovery of hypercalcaemia, and her postmenopausal status was confirmed by the finding of raised gonadotrophin concentrations (follicle stimulating hormone 93 U/l, luteinising hormone 109 U/l).

### Case 3

A 51 year old woman had been treated with vitamin D analogues and thyroxine supplements for 13 years since developing persistent hypoparathyroidism and hypothyroidism after subtotal thyroidectomy for Graves' disease. Her serum calcium concentrations had been within the reference range until October 1988, when she presented with a 6 month history of weight loss of 9 kg, nausea, vomiting, thirst, and intermittent confusion. She was also menopausal. There had been no change in her dose of 1 $\alpha$ -hydroxycholecalciferol (2 g/day), calcium intake, or compliance with treatment. Hypercalcaemia was confirmed (the calcium concentration, adjusted for albumin, was 3.25 mmol/l). Her calcium concentration returned to normal and her symptoms resolved after intravenous rehydration treatment and an interval without vitamin D therapy. Treatment with 1 $\alpha$ -hydroxycholecalciferol was reintroduced at a reduced daily dose of 0.25 g, and the patient's calcium concentrations remained within the reference range thereafter.

### Comment

These three cases show that a change in oestrogen status can alter sensitivity to a potent vitamin D analogue in women who do not have the ability to produce parathyroid hormone. In a similar case, reported in 1979, a patient became hypercalcaemic after stopping the oral contraceptive pill.<sup>2,3</sup> Reintroduction of oestrogen was associated with a fall in her serum calcium concentration. The anti-oestrogenic activity of danazol—prescribed for endometriosis in a patient with idiopathic hypoparathyroidism who was being treated with 1 $\alpha$ -hydroxycholecalciferol—resulted in hypercalcaemia and a reduced maintenance requirement for 1 $\alpha$ -hydroxycholecalciferol.<sup>4</sup> Hypercalcaemia can also occur immediately after delivery in women with hypoparathyroidism treated with vitamin D supplements.<sup>5,6</sup> All these observations support a crucial role for oestrogen in calcium regulation in these women.

Oestrogen, 1,25-dihydroxyvitamin D, and parathyroid hormone influence bone metabolism. Cytokines are now recognised as pivotal mediators of oestrogen, which acts on oestrogen receptors on osteoblasts and osteoclasts to inhibit bone resorption.<sup>9</sup> In normal women, oestrogen withdrawal increases bone resorption and causes a rise in serum calcium. 1,25-dihydroxyvitamin D is now known to be the major direct regulator of active transcellular calcium absorption via vitamin D receptors in intestinal mucosal cells.<sup>10</sup> Oestrogen can increase calcium absorption directly and indirectly by stimulating 1 $\alpha$ -hydroxylase activity in the kidney.<sup>11</sup> Withdrawal of oestrogens would theoretically reduce calcium absorption, and hypercalcaemia in these cases cannot be explained by this

mechanism. Ultimately, the effects of oestrogen on bone and calcium metabolism are monitored by the calcium sensing receptor on parathyroid cells,<sup>12</sup> which respond by altering parathyroid hormone secretion. Parathyroid hormone is a major modulator of osteoclast activity. In the absence of parathyroid hormone, the positive effect on osteoclast activity and bone resorption of withdrawal of oestrogen becomes much more important for calcium regulation. In the absence of parathyroid hormone, the balance between the action of 1,25-dihydroxyvitamin D, which is a potent inducer of bone resorption,<sup>13</sup> and of oestrogen, which inhibits bone resorption, may become more crucial. A clinical awareness of this phenomenon allows appropriate monitoring of patients and adjustment of their dose of vitamin D at the menopause or while starting or stopping hormone replacement therapy.

Contributors: JM, RD and AA collected cases. JM, FD and JH wrote the manuscript and are guarantors for the paper.

- Gann DS, Paone JF. Delayed hypocalcemia after thyroidectomy for Graves' disease is prevented by parathyroid autotransplantation. *Ann Surg* 1979;190:508-13.
- Verbeelen D, Fuss M. Hypercalcaemia induced by oestrogen withdrawal in vitamin D-treated hypoparathyroidism. *BMJ* 1979;i:522-3.
- Nagant de Deuxchaisnes C. Oestrogen-induced hypocalcaemia in hypoparathyroidism. *BMJ* 1979;i:1563.
- Hepburn NC, Abdul-Aziz LAS, Whiteoak R. Danazol-induced hypercalcaemia in alfacalcidol-treated hypoparathyroidism. *Postgrad Med J* 1989;65:849-50.
- Cundy T, Haining SA, Guillard-Cumming DF, Butler J, Kanis JA. Remission of hypoparathyroidism during lactation: evidence for a physiological role for prolactin in the regulation of vitamin D metabolism. *Clin Endocrinol* 1987;26:667-74.
- Wright AD, Joplin GF, Dixon HG. Postpartum hypercalcaemia in treated hypoparathyroidism. *BMJ* 1969;i:23-5.
- Eriksen EF, Colvard DS, Berg NJ, Graham ML, Mann KG, Spelsberg TC, et al. Evidence of oestrogen receptors on normal human osteoblast-like cells. *Science* 1988;241:84-6.
- Oursler MJ, Pederson L, Pyfferoen J, Osdoby P, Fitzpatrick L, Spelsberg TC. Oestrogen modulation of avian osteoclast lysosomal gene expression. *Endocrinology* 1993;132:1373-80.
- Manolagas SC, Jilka RL. Bone marrow cytokines and bone remodelling. Emerging insights into the pathophysiology of osteoporosis. *N Engl J Med* 1995;332:305-11.
- Reichel H, Koeffler HP, Norman AW. Role of vitamin D endocrine system in health and disease. *N Engl J Med* 1989;320:980.
- Gennari C, Agnusdei D, Nardi P, Givirelli R. Estrogen preserves a normal intestinal responsiveness to 1,25-dihydroxyvitamin D3 in oophorectomized women. *J Clin Endocrinol Metab* 1990;71:1288-93.
- Brown EM, Gamba G, Riccardi D, Lombardi M, Butters R, Kifor O, et al. Cloning and characterisation of an extracellular calcium sensing receptor from bovine parathyroid. *Nature* 1993;366:575-80.
- Reynolds JJ, Pavlovitch H, Balsan S. 1,25-dihydroxycholecalciferol increases bone resorption on thyroparathyroidectomized mice. *Calcified Tiss Res* 1976;21:207-12.

(Accepted 12 May 1999)

### Corrections and clarifications

Screening of newborn infants for cholestatic hepatobiliary disease with tandem mass spectrometry  
In this paper by I Mushtaq et al (21 August, pp 471-7) the x axis in figure 3 should have been labelled 1 $\alpha$ -specificity.

### Minerva

Minerva has made a mistake—the fourth paragraph in the issue of 2 October reports a trial of cognitive behaviour therapy in depression. It was published in *Archives of General Psychiatry* 1999;56:829-35 (not the *British Journal of Psychiatry*).

### Letters

In the letter by Philip C Herbert (26 June, p 1762), a wrong currency conversion was given. \$C40 000 is equivalent to about £16 300 [not £92 000].

University  
Department of  
Medicine, Glasgow  
Royal Infirmary  
John Hinnie  
lecturer  
Rosemary Dargie  
clinical assistant  
Alaa Al-Rawi  
senior house officer

Correspondence to:  
J McIlroy  
JMcIlroy@gri  
biochem.org.uk